

Ghost Kinetic Operator of Vacuum String Field Theory

Kazumi Okuyama

Enrico Fermi Institute, University of Chicago
5640 S. Ellis Ave., Chicago IL 60637, USA
`kazumi@theory.uchicago.edu`

Using the data of eigenvalues and eigenvectors of Neumann matrices in the 3-string vertex, we prove analytically that the ghost kinetic operator of vacuum string field theory obtained by Hata and Kawano is equal to the ghost operator inserted at the open string midpoint. We also comment on the values of determinants appearing in the norm of sliver state.

January 2002

1. Introduction

Vacuum string field theory (VSFT) [1] is proposed as the theory around the closed string vacuum after the open string tachyon condensation. (See [2–20] for related papers.) Since all the information of tachyon condensation is contained in the kinetic operator Q of this theory, it is important to determine its form and study its properties. One possible form of Q is

$$Q = c_0 + \sum_{n=1}^{\infty} (c_n + (-1)^n c_{-n}) f_n \quad (1.1)$$

where f_n 's are numerical coefficients. This satisfies the requirement that Q has a trivial cohomology and it is a derivation of the star product.

In [9], Hata and Kawano found that the coefficients f_n are determined uniquely if we demand that the equation of motion has a nontrivial solution in the Siegel gauge. The coefficients f_n^{HK} they found are written in terms of the Neumann coefficients in the 3-string vertex of ghost sector [9,10]:

$$f^{\text{HK}} = \frac{1}{1 - \widetilde{M}} \widetilde{v}. \quad (1.2)$$

See section 2 for the definition of $\widetilde{M}$ and $\widetilde{v}$.

Recently, Gaiotto, Rastelli, Sen and Zwiebach [17] discussed that there is a canonical choice of kinetic operator: the ghost insertion at the open string midpoint

$$Q = c \left(\frac{\pi}{2} \right) = \sum_{n=-\infty}^{\infty} c_n \cos \frac{\pi n}{2} = c_0 + \sum_{n=1}^{\infty} (c_n + (-1)^n c_{-n}) \cos \frac{\pi n}{2}. \quad (1.3)$$

This operator has a form (1.1) with the coefficients f_n given by

$$f_n^{\text{mid}} = \cos \frac{\pi n}{2} = \frac{1}{2} [i^n + (-i)^n]. \quad (1.4)$$

They conjectured that the operator found in [9] is nothing but this canonical kinetic operator, *i.e.*,

$$f^{\text{HK}} = f^{\text{mid}}, \quad (1.5)$$

and analyzed this equality numerically.

In this paper, we will prove (1.5) analytically. To do that, detailed information of the spectrum of matrix $\widetilde{M}$ is needed. Since $\widetilde{M}$ can be written in terms of the Neumann matrix M in the matter sector, we can use the spectrum of M recently obtained in [19].

This paper is organized as follows: In section 2, we review the spectrum of Neumann matrices obtained in [19]. In section 3, we show that the eigenvector of Neumann matrix is δ -function normalizable. In section 4, we prove the equation $f^{\text{HK}} = f^{\text{mid}}$ by showing that their generating functions coincide. In section 5, we define the index of Neumann matrix and compute it for some examples. In section 6, we estimate the values of determinants which appear in the norm of sliver state. It is found that the norm of the matter part is vanishing and the norm of the ghost part is divergent.

2. Eigenvalues and Eigenvectors of Neumann Matrices

In this section, we review the result of [19]. The 3-string vertex in the zero-momentum sector is given by [21]

$$|V_3\rangle = \exp\left(\sum_{r,s=1}^3 -\frac{1}{2}a^{(r)\dagger}V^{rs}a^{(s)\dagger} + c^{(r)\dagger}\tilde{V}^{rs}b^{(s)\dagger} + c^{(r)\dagger}\tilde{v}^{rs}b_0^{(s)}\right) \bigotimes_{r=1}^3 c_0^{(r)}c_1^{(r)}|0\rangle_r. \quad (2.1)$$

In this paper, we use the following notation for Neumann matrices:

$$M = CV^{11}, \quad \tilde{M} = C\tilde{V}^{11}, \quad \tilde{v} = \tilde{v}^{11}, \quad (2.2)$$

where $C_{nm} = (-1)^n\delta_{nm}$ is the twist matrix. Note that M , $\tilde{M}$ and $\tilde{v}$ are twist-even:

$$[M, C] = [\tilde{M}, C] = 0, \quad C\tilde{v} = \tilde{v}. \quad (2.3)$$

Since $|V_3\rangle$ is invariant under the action of $L_1 + L_{-1}$, the matter Neumann matrix M and the matrix K_1 can be simultaneously diagonalized, where K_1 is defined as the representation matrix of $L_1 + L_{-1}$ on matter oscillators:

$$[L_1 + L_{-1}, v \cdot a] = (K_1 v) \cdot a - \sqrt{2}v_1 p. \quad (2.4)$$

Here $v \cdot a = \sum_{n=1}^{\infty} v_n a_n$. The explicit form of K_1 is given by

$$(K_1)_{nm} = -\sqrt{n(n-1)}\delta_{n-1,m} - \sqrt{n(n+1)}\delta_{n+1,m}. \quad (2.5)$$

Since K_1 is a real and symmetric matrix, its eigenvalue is a real number. The spectrum of K_1 is continuous on the real axis. The eigenvector $v^{(k)}$ of K_1 with eigenvalue k is implicitly given by the generating function

$$f_k(z) = \sum_{n=1}^{\infty} \frac{v_n^{(k)}}{\sqrt{n}} z^n = \frac{1}{k} (1 - e^{-k \tan^{-1} z}). \quad (2.6)$$

This function has a symmetry

$$f_k(-z) = -f_{-k}(z), \quad (2.7)$$

which reflects the fact that K_1 is twist-odd

$$\{K_1, C\} = 0. \quad (2.8)$$

From $f_{k=0}(z) = \tan^{-1} z$, the eigenvector $v^{(0)}$ with eigenvalue $k = 0$ becomes

$$v_{2l}^{(0)} = 0, \quad v_{2l-1}^{(0)} = \frac{(-1)^{l-1}}{\sqrt{2l-1}}. \quad (2.9)$$

It is convenient to introduce a bracket notation for the infinite summation in (2.6):

$$f_k(z) = \langle z | E^{-1} | k \rangle = \langle k | E^{-1} | z \rangle, \quad (2.10)$$

where E is a diagonal matrix defined by

$$E_{nm} = \sqrt{n} \delta_{n,m}, \quad (2.11)$$

and $|z\rangle$ and $|k\rangle$ denote the infinite-dimensional vectors

$$|z\rangle = (z, z^2, z^3, \dots)^T, \quad |k\rangle = (v_1^{(k)}, v_2^{(k)}, v_3^{(k)}, \dots)^T. \quad (2.12)$$

$\langle z| = |z\rangle^T$ is the transpose of $|z\rangle$, not the hermitian conjugate of $|z\rangle$. Under the twist, $|z\rangle$ and $|k\rangle$ transform as

$$C|z\rangle = |-z\rangle, \quad C|k\rangle = -|-k\rangle. \quad (2.13)$$

A useful relation satisfied by $\langle z|$ is

$$z \partial_z \langle z| = \langle z| E^2. \quad (2.14)$$

For a matrix X which commutes with K_1 , let $X(k)$ denote the eigenvalue for the eigenvector $|k\rangle$

$$X|k\rangle = X(k)|k\rangle. \quad (2.15)$$

The eigenvalue of M is given by

$$M(k) = -\frac{1}{1 + 2 \cosh \frac{\pi k}{2}}. \quad (2.16)$$

The width-matrix T_N of the wedge state [22], which is defined by

$$|N\rangle_w = (|0\rangle)_\star^{N-1} = \exp\left(-\frac{1}{2}a^\dagger C T_N a^\dagger\right) |0\rangle, \quad (2.17)$$

also commutes with K_1 . T_N can be written as [8]

$$T_N = \frac{T + (-T)^{N-1}}{1 - (-T)^N}, \quad (2.18)$$

where T is the width-matrix of the sliver [23,2]

$$T = \frac{1}{2M} \left(1 + M - \sqrt{(1-M)(1+3M)}\right). \quad (2.19)$$

Note that $T_3 = M$ and $T_\infty = T$. The eigenvalues of T and T_N are given by

$$T(k) = -e^{-\frac{\pi}{2}|k|}, \quad T_N(k) = \frac{\sinh\left(\frac{2-N}{4}\pi k\right)}{\sinh\left(\frac{N}{4}\pi k\right)}. \quad (2.20)$$

3. Inner Product $\langle k|p\rangle$

In this section, we study the inner product between two eigenvectors $|k\rangle, |p\rangle$ of K_1 . The inner product of two vectors is defined by

$$\langle v|v'\rangle \equiv \sum_{n=1}^{\infty} v_n v'_n = \int_{-\frac{\pi}{2}}^{\frac{3\pi}{2}} \frac{d\theta}{2\pi} \langle v|e^{i\theta}\rangle \langle e^{-i\theta}|v'\rangle. \quad (3.1)$$

Since K_1 is symmetric, we can expect that two eigenvectors with different eigenvalues are orthogonal to each other with respect to this inner product. But the norm of eigenvector is divergent, which can be seen from the example (2.9). Therefore, the eigenvector of K_1 is non-normalizable in this sense. However, as we will show below, the eigenvector of K_1 is δ -function normalizable as usual for the continuous spectrum. To see this, let us calculate the inner product $\langle k|p\rangle$ of two eigenvectors:

$$\begin{aligned} \langle k|p\rangle &= \int_{-\frac{\pi}{2}}^{\frac{3\pi}{2}} \frac{d\theta}{2\pi} \langle k|E^{-1}|e^{i\theta}\rangle \langle e^{-i\theta}|E|p\rangle \\ &= \int_{-\frac{\pi}{2}}^{\frac{3\pi}{2}} \frac{d\theta}{2\pi} \langle k|E^{-1}|e^{i\theta}\rangle \left(z \partial_z \langle z|E^{-1}|p\rangle \Big|_{z=e^{-i\theta}} \right) \\ &= \int_{-\frac{\pi}{2}}^{\frac{3\pi}{2}} \frac{d\theta}{2\pi} \frac{1}{k} \left(1 - e^{-k \tan^{-1} e^{i\theta}} \right) \left(\frac{1}{2 \cos \theta} e^{-p \tan^{-1} e^{-i\theta}} \right) \end{aligned} \quad (3.2)$$




Fig.1 Image of the unit circle under the map $f(z) = \tan^{-1} z$

Here we used (2.14), (2.10) and (2.6).

In order to evaluate this integral, we should specify the branch of the function

$$\tan^{-1} z = \frac{1}{2i} \log \frac{1+iz}{1-iz}. \quad (3.3)$$

For the consistency of the relation $f_k(0) = 0$ (2.6), we should take a branch such that $\tan^{-1}(0) = 0$. In other words, $\tan^{-1} z$ is given by the Taylor series

$$\tan^{-1} z = \sum_{l=1}^{\infty} \frac{(-1)^{l-1}}{2l-1} z^{2l-1}. \quad (3.4)$$

On this branch of the map $w = \tan^{-1} z$, the unit disk $|z| < 1$ is mapped to the strip $|\operatorname{Re} w| < \pi/4$, the right half of the unit circle (C_R) is mapped to the line $\frac{\pi}{4} + i\mathbf{R}$ on the w -plane, and the left half (C_L) is mapped to $-\frac{\pi}{4} + i\mathbf{R}$ (see Fig.1):

$$\begin{aligned} \tan^{-1} e^{i\theta} &= \frac{\pi}{4} + \frac{i}{2} \log \tan \left(\frac{\theta}{2} + \frac{\pi}{4} \right) \quad \text{on } C_R = \left\{ -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2} \right\}, \\ &= -\frac{\pi}{4} - \frac{i}{2} \log \tan \left(\frac{\theta}{2} - \frac{\pi}{4} \right) \quad \text{on } C_L = \left\{ \frac{\pi}{2} \leq \theta \leq \frac{3\pi}{2} \right\}. \end{aligned} \quad (3.5)$$

Let us first consider the contribution from C_R . We make a change of integration variable from θ to x defined by

$$\tan^{-1} e^{i\theta} = \frac{\pi}{4} + ix, \quad \tan^{-1} e^{-i\theta} = \frac{\pi}{4} - ix, \quad \tan \frac{\theta}{2} = \tanh x, \quad (3.6)$$

$$\frac{d\theta}{2\cos\theta} = dx, \quad d\theta = \frac{2dx}{\cosh 2x}. \quad (3.7)$$

Note that x runs from $-\infty$ to ∞ . Then, the contribution from C_R to the integral (3.2) becomes

$$\begin{aligned} \int_{C_R} \frac{d\theta}{2\pi} \frac{1}{k} \left(1 - e^{-k \tan^{-1} e^{i\theta}}\right) \left(\frac{1}{2\cos\theta} e^{-p \tan^{-1} e^{-i\theta}}\right) &= \int_{-\infty}^{\infty} \frac{dx}{2\pi} \frac{1}{k} (1 - e^{-\frac{\pi k}{4} - ikx}) e^{-\frac{\pi p}{4} + ipx} \\ &= \frac{1}{k} \delta(p) - \frac{1}{k} e^{-\frac{\pi k}{2}} \delta(k - p). \end{aligned} \quad (3.8)$$

Next we consider the integral over C_L . Here we also change the integration variable as

$$\tan^{-1} e^{i\theta} = -\frac{\pi}{4} - ix, \quad \tan^{-1} e^{-i\theta} = -\frac{\pi}{4} + ix, \quad -\cot \frac{\theta}{2} = \tanh x, \quad (3.9)$$

$$\frac{d\theta}{2\cos\theta} = -dx, \quad d\theta = \frac{2dx}{\cosh 2x}. \quad (3.10)$$

The contribution from C_L is given by

$$\begin{aligned} \int_{C_L} \frac{d\theta}{2\pi} \frac{1}{k} \left(1 - e^{-k \tan^{-1} e^{i\theta}}\right) \left(\frac{1}{2\cos\theta} e^{-p \tan^{-1} e^{-i\theta}}\right) &= - \int_{-\infty}^{\infty} \frac{dx}{2\pi} \frac{1}{k} (1 - e^{\frac{\pi k}{4} + ikx}) e^{\frac{\pi p}{4} - ipx} \\ &= -\frac{1}{k} \delta(p) + \frac{1}{k} e^{\frac{\pi k}{2}} \delta(k - p). \end{aligned} \quad (3.11)$$

By adding (3.8) and (3.11), we can see that $\langle k|p\rangle$ is proportional to the δ -function

$$\langle k|p\rangle = \mathcal{N}(k) \delta(k - p), \quad (3.12)$$

where $\mathcal{N}(k)$ is given by

$$\mathcal{N}(k) = \frac{2}{k} \sinh \frac{\pi k}{2}. \quad (3.13)$$

From (3.8) and (3.11), k can be regarded as an analogue of momentum along the lines $\pm \frac{\pi}{4} + i\mathbf{R}$ on the w -plane.

We can introduce the normalized eigenvector $|\widehat{k}\rangle$ of K_1 by

$$|\widehat{k}\rangle = \mathcal{N}(k)^{-\frac{1}{2}} |k\rangle, \quad (3.14)$$

whose inner product is given by the δ -function

$$\langle \widehat{k} | \widehat{p} \rangle = \delta(k - p). \quad (3.15)$$

In terms of this basis, we can write down the completeness relation

$$\mathbf{1} = \int_{-\infty}^{\infty} dk |\widehat{k}\rangle \langle \widehat{k}| = \int_{-\infty}^{\infty} dk \mathcal{N}(k)^{-1} |k\rangle \langle k|. \quad (3.16)$$

As a consistency check of (3.16), let us prove the following relation

$$\langle z|w\rangle = \int_{-\infty}^{\infty} dk \langle z|E|\widehat{k}\rangle \langle \widehat{k}|E^{-1}|w\rangle \quad (3.17)$$

for arbitrary points z, w on the unit disk. The left-hand-side is given by

$$\langle z|w\rangle = \sum_{n=1}^{\infty} z^n w^n = \frac{zw}{1-zw}. \quad (3.18)$$

The right-hand-side is calculated as

$$\begin{aligned} & \int_{-\infty}^{\infty} dk \langle z|E|\widehat{k}\rangle \langle \widehat{k}|E^{-1}|w\rangle \\ &= z \partial_z \int_{-\infty}^{\infty} dk \langle z|E^{-1}|k\rangle \mathcal{N}(k)^{-1} \langle k|E^{-1}|w\rangle \\ &= \frac{z}{1+z^2} \int_{-\infty}^{\infty} dk \frac{k}{2 \sinh \frac{\pi k}{2}} e^{-k \tan^{-1} z} \frac{1}{k} (1 - e^{-k \tan^{-1} w}) \\ &= \frac{z}{1+z^2} \int_0^{\infty} \frac{dk}{\sinh \frac{\pi k}{2}} \left[-\sinh(k \tan^{-1} z) + \sinh(k \tan^{-1} z + k \tan^{-1} w) \right] \\ &= \frac{z}{1+z^2} \left[-\tan(\tan^{-1} z) + \tan(\tan^{-1} z + \tan^{-1} w) \right] \\ &= \frac{zw}{1-zw}. \end{aligned} \quad (3.19)$$

Here we have used the formula

$$\int_0^{\infty} dx \frac{\sinh ax}{\sinh bx} = \frac{\pi}{2b} \tan \frac{\pi a}{2b}, \quad (|\operatorname{Re} a| < \operatorname{Re} b). \quad (3.20)$$

Since the unit disk $|z| < 1$ is mapped to the strip $|\operatorname{Re}(\tan^{-1} z)| < \pi/4$, the k -integral in (3.19) converges.

4. Proof of $f^{\text{HK}} = f^{\text{mid}}$

In this section, we prove the equivalence of f^{HK} and f^{mid} using the result in the previous section. We will show that the generating functions of f_n^{HK} and f_n^{mid} coincide:

$$\langle z|f^{\text{HK}}\rangle = \langle z|f^{\text{mid}}\rangle. \quad (4.1)$$

The generating function of f^{mid} is easily found to be

$$\langle z|f^{\text{mid}}\rangle = \sum_{n=1}^{\infty} z^n \cos \frac{\pi n}{2} = \sum_{l=1}^{\infty} (-1)^l z^{2l} = -\frac{z^2}{1+z^2}. \quad (4.2)$$

We assume $|z| < 1$ to make this summation converge. In order to calculate $\langle z|f^{\text{HK}}\rangle$, we have to know the explicit form of Neumann coefficients in the ghost sector. $\tilde{v}$ is given by [21]

$$\tilde{v}_{2l} = \frac{2}{3} B_{2l}, \quad \tilde{v}_{2l-1} = 0, \quad (4.3)$$

where B_{2l} is defined by

$$\left(\frac{1+iz}{1-iz} \right)^{\frac{2}{3}} = \exp \left(\frac{4i}{3} \tan^{-1} z \right) = \sum_{n=\text{even}} B_n z^n + i \sum_{n=\text{odd}} B_n z^n. \quad (4.4)$$

The generating function of $\tilde{v}$ becomes

$$\langle z|\tilde{v}\rangle = \frac{2}{3} \cosh \left(\frac{4}{3} i \tan^{-1} z \right). \quad (4.5)$$

The Neumann matrix $\widetilde{M}$ in the ghost sector is related to the corresponding matrix M in the matter sector as [21]

$$\widetilde{M} = -E \left(\frac{M}{1+2M} \right) E^{-1}. \quad (4.6)$$

Thus, f^{HK} can be written as

$$f^{\text{HK}} = \frac{1}{1-\widetilde{M}} \tilde{v} = E A E^{-1} \tilde{v}, \quad (4.7)$$

where

$$A = \frac{1+2M}{1+3M}. \quad (4.8)$$

Therefore, we can use the information of the spectrum of M to calculate the generating function $\langle z|f^{\text{HK}}\rangle$. Plugging $M(k)$ of the form (2.16) into (4.8), the eigenvalue of A is found to be

$$A(k) = \frac{2 \cosh \frac{\pi k}{2} - 1}{2 \cosh \frac{\pi k}{2} - 2}. \quad (4.9)$$

Using the completeness condition (3.16), $\langle z|f^{\text{HK}}\rangle$ can be rewritten as

$$\begin{aligned}
\langle z|f^{\text{HK}}\rangle &= \int_{-\infty}^{\infty} dk \langle z|E|k\rangle \langle k|E^{-1}|f^{\text{HK}}\rangle \\
&= z\partial_z \int_{-\infty}^{\infty} dk \langle z|E^{-1}|k\rangle \mathcal{N}(k)^{-1} \langle k|E^{-1}|f^{\text{HK}}\rangle \\
&= \frac{z}{1+z^2} \int_{-\infty}^{\infty} dk \frac{e^{-k \tan^{-1} z}}{2 \sinh \frac{\pi k}{2}} k \langle k|E^{-1}|f^{\text{HK}}\rangle \\
&= -\frac{z}{1+z^2} \int_0^{\infty} dk \frac{\sinh(k \tan^{-1} z)}{\sinh \frac{\pi k}{2}} k \langle k|E^{-1}|f^{\text{HK}}\rangle.
\end{aligned} \tag{4.10}$$

In the last step, we used the fact that $\langle k|E^{-1}|f^{\text{HK}}\rangle$ is an odd-function of k which follows from $C|k\rangle = -|-k\rangle$ and $C|f^{\text{HK}}\rangle = |f^{\text{HK}}\rangle$:

$$\langle -k|E^{-1}|f^{\text{HK}}\rangle = -\langle k|CE^{-1}|f^{\text{HK}}\rangle = -\langle k|E^{-1}C|f^{\text{HK}}\rangle = -\langle k|E^{-1}|f^{\text{HK}}\rangle. \tag{4.11}$$

From (4.7), the last factor in (4.10) is written as

$$k \langle k|E^{-1}|f^{\text{HK}}\rangle = A(k)k \langle k|E^{-1}|\tilde{v}\rangle. \tag{4.12}$$

Therefore, the problem is reduced to the calculation of $k \langle k|E^{-1}|\tilde{v}\rangle$. This quantity can be extracted from the generating function of $\tilde{v}$ (4.5):

$$\begin{aligned}
k \langle k|E^{-1}|\tilde{v}\rangle &= \int_{-\frac{\pi}{2}}^{\frac{3\pi}{2}} \frac{d\theta}{2\pi} k \langle k|E^{-1}|e^{i\theta}\rangle \langle e^{-i\theta}|\tilde{v}\rangle \\
&= \int_{-\frac{\pi}{2}}^{\frac{3\pi}{2}} \frac{d\theta}{2\pi} (1 - e^{-k \tan^{-1} e^{i\theta}}) \frac{2}{3} \cosh \left(\frac{4}{3} i \tan^{-1} e^{-i\theta} \right).
\end{aligned} \tag{4.13}$$

By changing the integration variable as in the previous section, this integral can be written as

$$\begin{aligned}
k \langle k|E^{-1}|\tilde{v}\rangle &= \frac{8}{3} \int_{-\infty}^{\infty} \frac{dx}{2\pi} \frac{1}{\cosh 2x} \left[1 - \cosh \left(\frac{\pi k}{4} + ikx \right) \right] \cosh \left(\frac{4}{3}x + \frac{\pi i}{3} \right) \\
&= \frac{8}{3} \int_0^{\infty} \frac{dx}{2\pi} \frac{1}{\cosh 2x} \left[\cosh \left(\frac{4}{3}x \right) - \cosh \left(\frac{4}{3}x + ikx \right) \cosh \left(\frac{\pi k}{4} + \frac{\pi i}{3} \right) \right. \\
&\quad \left. - \cosh \left(\frac{4}{3}x - ikx \right) \cosh \left(\frac{\pi k}{4} - \frac{\pi i}{3} \right) \right].
\end{aligned} \tag{4.14}$$

By making use of the formula

$$\int_0^{\infty} dx \frac{\cosh ax}{\cosh bx} = \frac{\pi}{2b} \cdot \frac{1}{\cos \frac{\pi a}{2b}}, \quad (|\text{Re } a| < \text{Re } b), \tag{4.15}$$

the integral in (4.14) is evaluated as

$$k\langle k|E^{-1}|\tilde{v}\rangle = \frac{1}{3} \left[2 - \frac{\cosh\left(\frac{\pi k}{4} + \frac{\pi i}{3}\right)}{\cosh\left(\frac{\pi k}{4} - \frac{\pi i}{3}\right)} - \frac{\cosh\left(\frac{\pi k}{4} - \frac{\pi i}{3}\right)}{\cosh\left(\frac{\pi k}{4} + \frac{\pi i}{3}\right)} \right] = \frac{2 \cosh \frac{\pi k}{2} - 2}{2 \cosh \frac{\pi k}{2} - 1} = \frac{1}{A(k)}. \quad (4.16)$$

Therefore,

$$k\langle k|E^{-1}|f^{\text{HK}}\rangle = kA(k)\langle k|E^{-1}|\tilde{v}\rangle = 1. \quad (4.17)$$

Finally, (4.10) becomes

$$\begin{aligned} \langle z|f^{\text{HK}}\rangle &= -\frac{z}{1+z^2} \int_0^\infty dk \frac{\sinh(k \tan^{-1} z)}{\sinh \frac{\pi k}{2}} \\ &= -\frac{z}{1+z^2} \tan(\tan^{-1} z) = -\frac{z^2}{1+z^2} = \langle z|f^{\text{mid}}\rangle. \end{aligned} \quad (4.18)$$

Here we used the formula (3.20). This is valid since $|\text{Re}(\tan^{-1} z)| < \pi/4$ for $|z| < 1$. This completes the proof of (4.1).

We comment on the subtlety of (4.17). Naively, one might think that the following equation holds:

$$k\langle k|E^{-1}|f^{\text{mid}}\rangle = k\langle k|E^{-1}|f^{\text{HK}}\rangle. \quad (4.19)$$

However, the left-hand-side is ill-defined since $\tan^{-1}(i) = i\infty$:

$$k\langle k|E^{-1}|f^{\text{mid}}\rangle = \frac{k}{2} [f_k(i) + f_k(-i)] = 1 - \cosh(k \tan^{-1}(i)). \quad (4.20)$$

By regularizing the last term of (4.20) with a parameter $a(< 1)$

$$k\langle k|E^{-1}|f^{\text{mid}}\rangle = \lim_{a \rightarrow 1} \frac{k}{2} [f_k(ia) + f_k(-ia)] = 1 - \lim_{a \rightarrow 1} \cosh(k \tan^{-1}(ia)), \quad (4.21)$$

we can show that this term does not contribute to the computation of $\langle z|f\rangle$:

$$\begin{aligned} &\lim_{a \rightarrow 1} \int_0^\infty dk \frac{\sinh(k \tan^{-1} z)}{\sinh \frac{\pi k}{2}} \cosh(k \tan^{-1}(ia)) \\ &= \lim_{a \rightarrow 1} \frac{1}{2} \int_0^\infty dk \frac{1}{\sinh \frac{\pi k}{2}} \left[\sinh(k \tan^{-1} z + k \tan^{-1}(ia)) + \sinh(k \tan^{-1} z - k \tan^{-1}(ia)) \right] \\ &= \lim_{a \rightarrow 1} \frac{1}{2} \left[\frac{z + ia}{1 - iaz} + \frac{z - ia}{1 + iaz} \right] = \lim_{a \rightarrow 1} \frac{(1 - a^2)z}{1 + a^2 z^2} = 0. \end{aligned} \quad (4.22)$$

Therefore, (4.19) should be understood as an equation up to such an irrelevant term. Here we emphasize that $\langle z|f^{\text{mid}}\rangle$ is well-defined and is equal to $\langle z|f^{\text{HK}}\rangle$ without any ambiguity.

5. Index of Neumann Matrix

As an application of our formalism, we define the index of Neumann matrix and compute it for some examples. For a general matrix X diagonalized on the basis $\{|k\rangle\}$ with its eigenvalue $X(k) = X(-k)$ an even-function of k , the eigenvalue $X(k) = X(-k)$ has a two-fold degeneracy

$$\begin{aligned} |k, +\rangle &= |k\rangle + C|k\rangle = |k\rangle - |-k\rangle, \\ |k, -\rangle &= |k\rangle - C|k\rangle = |k\rangle + |-k\rangle, \end{aligned} \quad (5.1)$$

where $|k, +\rangle$ ($|k, -\rangle$) is a twist-even (odd) eigenvector. Note that X is twist-even $[X, C] = 0$ when $X(k) = X(-k)$. For $k = 0$, there is only one twist-odd eigenvector. Therefore, the index of X defined by

$$I(X) = \text{Tr}(CX) \quad (5.2)$$

receives contribution only from $k = 0$. This can be thought of as a measure of the twist anomaly [13]. $I(X)$ can be calculated as

$$\begin{aligned} I(X) &= \int_{-\infty}^{\infty} dk \langle \hat{k} | CX | \hat{k} \rangle = \int_{-\infty}^{\infty} dk \langle \hat{k} | C | \hat{k} \rangle X(k) \\ &= - \int_{-\infty}^{\infty} dk \langle \hat{k} | - \hat{k} \rangle X(k) = - \int_{-\infty}^{\infty} dk \delta(2k) X(k) = -\frac{1}{2} X(0). \end{aligned} \quad (5.3)$$

As expected, it depends only on the eigenvalue at $k = 0$.

For example, the index of T_N is given by

$$I(T_N) = -\frac{1}{2} T_N(0) = \frac{1}{2} - \frac{1}{N}. \quad (5.4)$$

Here we used (2.20). In the rest of this section, we check this prediction for some examples by directly computing the trace. We first consider the index of B defined by [19]

$$B = \frac{d}{dN} T_N \Big|_{N=2} = -\frac{T \log(-T)}{1 - T^2}. \quad (5.5)$$

The eigenvalue of B is given by

$$B(k) = -\frac{1}{2} \frac{\frac{\pi k}{2}}{\sinh \frac{\pi k}{2}}. \quad (5.6)$$

It is known that this matrix can be written explicitly in the n, m basis [19]

$$B_{nm} = -\frac{1 + (-1)^{n+m}}{2} \frac{(-1)^{\frac{n-m}{2}} \sqrt{nm}}{(n+m)^2 - 1}. \quad (5.7)$$

Therefore, we can compute the index of B directly as

$$I(B) = \sum_{n=1}^{\infty} (-1)^n B_{nn} = - \sum_{n=1}^{\infty} (-1)^n \frac{n}{(2n)^2 - 1} = \frac{1}{4} \sum_{n=1}^{\infty} \left(\frac{(-1)^{n-1}}{2n-1} - \frac{(-1)^n}{2n+1} \right) = \frac{1}{4}. \quad (5.8)$$

This agrees with the expected result

$$I(B) = -\frac{1}{2}B(0) = \frac{d}{dN}I(T_N)\Big|_{N=2} = \frac{1}{4}. \quad (5.9)$$

Next we consider the index of $T_3 = M$. The diagonal element of M is given by [21]

$$M_{nn} = -\frac{1}{3} \left[2 \sum_{k=0}^n (-1)^k A_k^2 - 1 - (-1)^n A_n^2 \right] \quad (5.10)$$

where A_k is defined by

$$\sum_{k=\text{even}} A_k z^k + i \sum_{k=\text{odd}} A_k z^k = \left(\frac{1+iz}{1-iz} \right)^{\frac{1}{3}} = \exp \left(\frac{2i}{3} \tan^{-1} z \right). \quad (5.11)$$

The partial sum of the series $(-1)^n M_{nn}$ turns out to be

$$\begin{aligned} \sum_{n=1}^L (-1)^n M_{nn} &= \frac{1}{3} \left(1 - \sum_{k=0}^L (-1)^k A_k^2 \right) \quad L = \text{even}, \\ &= \frac{1}{3} \sum_{k=0}^L (-1)^k A_k^2 \quad L = \text{odd}. \end{aligned} \quad (5.12)$$

This sum converges in the limit $L \rightarrow \infty$ since $\sum_{k=0}^{\infty} (-1)^k A_k^2 = \frac{1}{2}$. This can be shown by integrating the generating function of A_k (5.11):

$$\begin{aligned} \sum_{k=0}^{\infty} (-1)^k A_k^2 &= \int_{-\frac{\pi}{2}}^{\frac{3\pi}{2}} \frac{d\theta}{2\pi} e^{\frac{2i}{3} \tan^{-1} e^{i\theta}} e^{\frac{2i}{3} \tan^{-1} e^{-i\theta}} \\ &= \int_{C_R} \frac{d\theta}{2\pi} e^{\frac{\pi i}{3}} + \int_{C_L} \frac{d\theta}{2\pi} e^{-\frac{\pi i}{3}} = \frac{1}{2}. \end{aligned} \quad (5.13)$$

Therefore, the index of M computed in this way agrees with the expected result (5.4)

$$I(M) = \sum_{n=1}^{\infty} (-1)^n M_{nn} = \frac{1}{6}. \quad (5.14)$$

It will be interesting to check (5.4) for general N by directly computing the trace.

We comment on a deformation of the index. We can introduce a parameter β in the definition of index, *e.g.*,

$$I_{\beta}(X) = \text{Tr}(CX e^{-\beta K_1^2}). \quad (5.15)$$

Naively, $I_{\beta}(X)$ is independent of β . However, it can be β -dependent as in the case of Witten index of a theory with continuous spectrum. It is interesting to study the β -(in)dependence of $I_{\beta}(X)$.

6. Norm of Sliver

6.1. Eigenvalue Density $\rho(k)$

In contrast with the finiteness of the index (5.2), the trace of X itself is divergent in general. This can be seen as

$$\text{Tr} X = \int_{-\infty}^{\infty} dk \langle \widehat{k} | X | \widehat{k} \rangle = \int_{-\infty}^{\infty} dk \langle \widehat{k} | \widehat{k} \rangle X(k) = \int_{-\infty}^{\infty} dk \delta(0) X(k). \quad (6.1)$$

If we formally define the eigenvalue density $\rho(k)$ by

$$\text{Tr} X = \int_{-\infty}^{\infty} dk \rho(k) X(k), \quad (6.2)$$

then $\rho(k)$ is independent of k and divergent

$$\rho(k) = \langle \widehat{k} | \widehat{k} \rangle = \frac{\langle k | k \rangle}{\mathcal{N}(k)} = \delta(0). \quad (6.3)$$

In the level truncation approximation, we can estimate the order of divergence of $\rho(k) = \rho(0)$ as follows:

$$\rho(k) = \frac{\langle k=0 | k=0 \rangle}{\mathcal{N}(0)} \sim \frac{1}{\mathcal{N}(0)} \sum_{n=1}^L \left(v_n^{(0)} \right)^2 \sim \frac{1}{\pi} \sum_{l=1}^{L/2} \frac{1}{2l-1} \sim \frac{1}{2\pi} \log L \quad (\text{for large } L). \quad (6.4)$$

This expression agrees with the one obtained in [19]. We can also estimate $\rho(k)$ by computing the trace of B in two different ways:

$$\text{Tr} B = \int_{-\infty}^{\infty} dk \rho(k) B(k) \sim \sum_{n=1}^L B_{nn} \quad (6.5)$$

where

$$\begin{aligned} \int_{-\infty}^{\infty} dk \rho(k) B(k) &= \rho(k) \int_{-\infty}^{\infty} dk - \frac{1}{2} \frac{\frac{\pi k}{2}}{\sinh \frac{\pi k}{2}} = -\frac{\pi}{2} \rho(k), \\ \sum_{n=1}^L B_{nn} &= -\sum_{n=1}^L \frac{n}{4n^2-1} \sim -\frac{1}{4} \log L. \end{aligned} \quad (6.6)$$

This again reproduce (6.4).

Now it is natural to define the regularized trace Tr_L by

$$\text{Tr}_L X = \log L \int_{-\infty}^{\infty} \frac{dk}{2\pi} X(k). \quad (6.7)$$

Using this trace, the determinant of X can be regularized as

$$\det X \sim \exp(\text{Tr}_L \log X) = L^{\gamma(X)}, \quad (6.8)$$

where the exponent $\gamma(X)$ is defined by

$$\gamma(X) = \int_{-\infty}^{\infty} \frac{dk}{2\pi} \log X(k). \quad (6.9)$$

When the eigenvalue $X(k)$ is an even-function of k , $\gamma(X)$ can be written as

$$\gamma(X) = \int_0^{\infty} \frac{dk}{\pi} \log X(k) = \frac{2}{\pi^2} \int_0^1 \frac{du}{u} \log X(k) \Big|_{u=e^{-\frac{\pi}{2}k}}. \quad (6.10)$$

6.2. Classical Solution of VSFT

In this subsection, we review the classical solution Ψ_0 of VSFT found in [9]. Ψ_0 is given by

$$|\Psi_0\rangle = -\mathcal{N}_m \mathcal{N}_g \exp\left(-\frac{1}{2}a^\dagger C T a^\dagger + c^\dagger C \tilde{T} b^\dagger\right) c_1 |0\rangle. \quad (6.11)$$

Here T is defined by (2.19) and $\tilde{T}$ is defined by

$$\tilde{T} = \frac{1}{2\widetilde{M}} \left(1 + \widetilde{M} - \sqrt{(1 - \widetilde{M})(1 + 3\widetilde{M})}\right) = -E T E^{-1}. \quad (6.12)$$

The matter part of Ψ_0 is the sliver state and the ghost part is conjectured to be the twisted sliver state [17]. $\mathcal{N}_m$ and $\mathcal{N}_g$ are the normalization constants given by

$$\begin{aligned} \mathcal{N}_m &= \det^{\frac{D}{2}}(1 - M)(1 + T), \\ \mathcal{N}_g &= \det^{-1}(1 - \widetilde{M})(1 + \tilde{T}), \end{aligned} \quad (6.13)$$

where $D(= 26)$ is the dimension of D-brane. The energy density of this solution is proportional to

$$\frac{(2\pi)^D}{V_D} \langle \Psi_0 | Q | \Psi_0 \rangle = E_m E_g, \quad (6.14)$$

where V_D is the volume of D-brane and

$$\begin{aligned} E_m &= \det(1 - M)^{\frac{3}{4}D} (1 + 3M)^{\frac{1}{4}D}, \\ E_g &= \det(1 - \widetilde{M})^{-\frac{3}{2}} \det(1 + 3\widetilde{M})^{-\frac{1}{2}}. \end{aligned} \quad (6.15)$$

6.3. Calculation of Various Determinants

We can estimate the values of determinants in (6.13) and (6.15) by using the regularization (6.8). Let us compute the exponent γ (6.10) for various matrices. One can show that γ 's of Neumann matrices are written as combinations of the following integrals:

$$\begin{aligned}
I_- &= \int_0^1 \frac{du}{u} \log(1-u) = - \int_0^1 \frac{du}{u} \sum_{n=1}^{\infty} \frac{1}{n} u^n = -S_0 = -\frac{\pi^2}{6}, \\
I_+ &= \int_0^1 \frac{du}{u} \log(1+u) = \int_0^1 \frac{du}{u} \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} u^n = -S_1 = \frac{\pi^2}{12}, \\
J_- &= \int_0^1 \frac{du}{u} \log(1-u+u^2) = \int_0^1 \frac{du}{u} \log(1-e^{\frac{\pi}{3}i}u)(1-e^{-\frac{\pi}{3}i}u) \\
&= - \int_0^1 \frac{du}{u} \sum_{n=1}^{\infty} \frac{1}{n} u^n (e^{\frac{\pi n}{3}i} + e^{-\frac{\pi n}{3}i}) = -2S_{\frac{1}{3}} = -\frac{\pi^2}{18}, \\
J_+ &= \int_0^1 \frac{du}{u} \log(1+u+u^2) = \int_0^1 \frac{du}{u} \log(1-e^{\frac{2\pi}{3}i}u)(1-e^{-\frac{2\pi}{3}i}u) \\
&= - \int_0^1 \frac{du}{u} \sum_{n=1}^{\infty} \frac{1}{n} u^n (e^{\frac{2\pi n}{3}i} + e^{-\frac{2\pi n}{3}i}) = -2S_{\frac{2}{3}} = \frac{\pi^2}{9},
\end{aligned} \tag{6.16}$$

where S_a is defined by

$$S_a \equiv \sum_{n=1}^{\infty} \frac{1}{n^2} \cos(\pi n a) = \frac{\pi^2}{4} (1-a)^2 - \frac{\pi^2}{12}, \quad (0 \leq a \leq 2). \tag{6.17}$$

Substituting the eigenvalues (2.16) and (2.20) into the definition of γ (6.10), the exponents of the matter Neumann matrices are found to be

$$\begin{aligned}
\gamma(1+T) &= \frac{2}{\pi^2} I_- = -\frac{1}{3}, & \gamma(1-T) &= \frac{2}{\pi^2} I_+ = \frac{1}{6} \\
\gamma(1-M) &= \frac{2}{\pi^2} (2I_+ - J_+) = \frac{1}{9}, & \gamma(1+3M) &= \frac{2}{\pi^2} (2I_- - J_+) = -\frac{8}{9} \\
\gamma(1+2M) &= \frac{2}{\pi^2} (J_- - J_+) = -\frac{1}{3}, & \gamma(1-TM) &= \frac{2}{\pi^2} (I_+ - J_+) = -\frac{1}{18}.
\end{aligned} \tag{6.18}$$

For the ghost part, using the relation

$$\tilde{T} = -ETE^{-1}, \quad 1 - \tilde{M} = E \left(\frac{1+3M}{1+2M} \right) E^{-1}, \quad 1 + 3\tilde{M} = E \left(\frac{1-M}{1+2M} \right) E^{-1}, \tag{6.19}$$

and assuming the property

$$\det(EXE^{-1}) = \det X, \tag{6.20}$$

the exponents can be computed as

$$\begin{aligned}
\gamma(1 + \widetilde{T}) &= \gamma(1 - T) = \frac{1}{6}, \\
\gamma(1 - \widetilde{M}) &= \gamma(1 + 3M) - \gamma(1 + 2M) = -\frac{5}{9}, \\
\gamma(1 + 3\widetilde{M}) &= \gamma(1 - M) - \gamma(1 + 2M) = \frac{4}{9}.
\end{aligned} \tag{6.21}$$

Combining these relations, the order of determinants (6.13) and (6.15) can be estimated:

$$\begin{aligned}
\mathcal{N}_m &\sim L^{-\frac{1}{9}D}, \quad \mathcal{N}_g \sim L^{\frac{7}{18}}, \\
E_m &\sim L^{-\frac{5}{36}D}, \quad E_g \sim L^{\frac{11}{18}}.
\end{aligned} \tag{6.22}$$

In the limit $L \rightarrow \infty$, $\mathcal{N}_m$ and E_m tend to vanish. This behavior of the norm in the matter sector was observed in the numerical calculation [2]. On the other hand, the norm in the ghost sector diverges in the limit $L \rightarrow \infty$. In the early days of the study of VSFT, it was expected that the divergent factor coming from the ghost sector compensates the vanishing factor from the matter sector and in total the norm becomes finite [2]. However, our result (6.22) shows that the contribution from the ghost sector is not large enough to compensate the matter contribution. The total norm is still vanishing in the limit $L \rightarrow \infty$.

Recent discussion in [17] is that the action with the kinetic operator (1.1) is a singular description of the theory around the closed string vacuum and there is an infinite overall factor in front of the action due to a singular field redefinition. Therefore, this infinite factor would cancel the vanishing norm of the sliver as discussed in [17].

Acknowledgement

I would like to thank Kentaro Hori and Sunny Itzhaki for discussion and encouragement.

References

- [1] L. Rastelli, A. Sen and B. Zwiebach, “String Field Theory Around the Tachyon Vacuum,” [hep-th/0012251]; “Vacuum string field theory,” [hep-th/0106010].
- [2] L. Rastelli, A. Sen and B. Zwiebach, “Classical Solutions in String Field Theory Around the Tachyon Vacuum,” [hep-th/0102112];
- [3] L. Rastelli, A. Sen and B. Zwiebach, “Half-strings, Projectors, and Multiple D-branes in Vacuum String Field Theory,” [hep-th/0105058]; “Boundary CFT construction of D-branes in vacuum string field theory,” [hep-th/0105168];
- [4] D. J. Gross and W. Taylor, “Split string field theory I,” JHEP **0108**, 009 (2001), [hep-th/0105059]; “Split string field theory. II,” JHEP **0108**, 010 (2001) [hep-th/0106036].
- [5] T. Kawano and K. Okuyama, “Open string fields as matrices,” JHEP **0106**, 061 (2001) [hep-th/0105129].
- [6] Y. Matsuo, “BCFT and sliver state,” Phys. Lett. B **513**, 195 (2001) [hep-th/0105175]; “Identity projector and D-brane in string field theory,” Phys. Lett. B **514**, 407 (2001) [hep-th/0106027]; “Projection operators and D-branes in purely cubic open string field theory,” Mod. Phys. Lett. A **16**, 1811 (2001) [hep-th/0107007].
- [7] J. R. David, “Excitations on wedge states and on the sliver,” JHEP **0107**, 024 (2001) [hep-th/0105184].
- [8] K. Furuuchi and K. Okuyama, “Comma vertex and string field algebra,” JHEP **0109**, 035 (2001) [hep-th/0107101].
- [9] H. Hata and T. Kawano, “Open string states around a classical solution in vacuum string field theory,” JHEP **0111**, 038 (2001) [hep-th/0108150].
- [10] I. Kishimoto, “Some properties of string field algebra,” JHEP **0112**, 007 (2001) [hep-th/0110124].
- [11] P. Mukhopadhyay, “Oscillator representation of the BCFT construction of D-branes in vacuum string field theory,” [hep-th/0110136].
- [12] N. Moeller, “Some exact results on the matter star-product in the half-string formalism,” [hep-th/0110204].
- [13] H. Hata and S. Moriyama, “Observables as Twist Anomaly in Vacuum String Field Theory,” [hep-th/0111034].
- [14] G. Moore and W. Taylor, “The singular geometry of the sliver,” [hep-th/0111069].
- [15] K. Okuyama, “Siegel gauge in vacuum string field theory,” [hep-th/0111087].
- [16] A. Hashimoto and N. Itzhaki, “Observables of string field theory,” [hep-th/0111092].
- [17] D. Gaiotto, L. Rastelli, A. Sen and B. Zwiebach, “Ghost Structure and Closed Strings in Vacuum String Field Theory,” [hep-th/0111129].
- [18] L. Rastelli, A. Sen and B. Zwiebach, “A note on a proposal for the tachyon state in vacuum string field theory,” [hep-th/0111153];

- R. Rashkov and K. S. Viswanathan, “A Note on the Tachyon State in Vacuum String Field Theory,” [hep-th/0112202].
- [19] L. Rastelli, A. Sen and B. Zwiebach, “Star algebra spectroscopy,” [hep-th/0111281].
 - [20] I. Kishimoto and K. Ohmori, “CFT description of identity string field: Toward derivation of the VSFT action,” [hep-th/0112169].
 - [21] D. J. Gross and A. Jevicki, “Operator Formulation of Interacting String Field Theory,” Nucl. Phys. **B283** (1987) 1; “Operator Formulation of Interacting String Field Theory (II),” Nucl. Phys. **B287** (1987) 225.
 - [22] L. Rastelli and B. Zwiebach, “Tachyon potentials, star products and universality,” JHEP **0109**, 038 (2001) [hep-th/0006240].
 - [23] V. A. Kostelecky and R. Potting, “Analytical construction of a nonperturbative vacuum for the open bosonic string,” Phys. Rev. **D63** (2001) 046007 [hep-th/0008252].